

Tidal disruption and the tale of three clusters

Guido De Marchi¹, Luigi Pulone², and Francesco Paresce³

¹ ESA, Space Science Department, Noordwijk, Netherlands gdemarchi@esa.int

² INAF, Observatory of Rome, Monte Porzio, Italy pulone@mporzio.astro.it

³ INAF, Rome, Italy fparesce@inaf.it

Abstract⁴

How well can we tell whether a cluster will survive the Galaxy’s tidal forces? This is conceptually easy to do if we know the cluster’s total mass, mass structure and space motion parameters. This information is used in models that predict the probability of disruption due to tidal stripping, disc and bulge shocking [7, 6, 2]. But just how accurate is the information that goes into these models and, therefore, how reliable are their predictions? To understand the virtues and weaknesses of these models, we have studied in detail three clusters (NGC 6397, NGC 6712, NGC 6218) whose predicted interaction with the galaxy is very different. We have used deep HST and VLT data to measure the luminosity function (LF) of stars throughout the clusters in order to derive a solid global mass function (GMF). The latter is the best tell-tale of the strength and extent of tidal stripping operated by the Galaxy. This is because the evaporation of stars from the cluster causes a progressive flattening of the IMF [9] and this effect is enhanced by tidal stripping and disc/bulge shocking. Therefore, at any time, the shape of the GMF must reflect the past interaction of the cluster with the Galaxy. Since the three clusters that we have studied have widely different probabilities of disruption (see the predicted times to disruption T_d in Table 1), we expected to find widely different GMFs. We indeed found that the GMF of the three clusters is different, but not in the way predicted by the models, as we explain below.

To derive a reliable GMF, it is necessary to measure the stellar LF at various locations in the cluster, in order to build a solid model of the cluster’s dynamics. Our accurate measurements in several bands [4, 1, 5] have allowed us to determine the GMF of the three clusters for stars down to $\sim 0.2 M_\odot$, where it is most sensitive to tidal stripping. While the GMF of NGC 6397 is that typical of globular clusters, with a peak at $\sim 0.3 M_\odot$, NGC 6218 and NGC 6712 have very flat GMF. Near the half-mass radius, the LF that we observed for these two clusters is severely depleted at the low-mass end. But while the orbit of NGC 6712 is compatible with “ferocious” stripping, that

⁴ Space limitations do not allow us to present our work in full detail. We, therefore, include here a brief summary of the main results. For the complete version of this paper, the reader should refer to De Marchi et al. (2006) [5].

of NGC 6218 is not [3]. However, our analysis shows that the orbit of NGC 6218 used in these models is not realistic. In fact, more recent work based on the Hipparcos reference system [8] indicates for NGC 6218 an irregular orbit with shorter perigalactic distance, more prone to extensive stripping. We conclude that existing models of cluster disruption are limited by the lack of information on the precise cluster space motions. However, they benefit from an accurate knowledge of the GMF. Before the advent of GAIA, which will make it possible to measure reliably the orbits of many clusters, the only way forward to understand the evolution of the globular cluster system rests on the accurate measurement of the GMF of a large number of these objects. A telescope such as the VLT is perfectly suited for this task.

Table 1. Predicted times to disruption T_d in Gyr

Reference	NGC 6397	NGC 6218	NGC 6712
Gnedin & Ostriker 1997	2.1	14.5	0.3
Dinescu et al. 1999	3.9	29.4	3.7
Baumgardt & Makino 2003	11.3	16.3	9.0




Fig. 1. The solid lines show the best fitting power-law GMF as derived using our dynamical models applied to a set of radial LFs. The symbols joined by the dashed lines show the local MF as measured near the half-mass radius of each cluster, proving that the latter is a good approximation to the GMF. While the time to disruption (T_d) of NGC 6712 justifies its falling GMF, NGC 6218 should have a GMF as steep as that of NGC 6397, given its very long T_d .

References

1. G. Andreuzzi, G. De Marchi, F. Ferraro, F. Paresce, L. Pulone, R. Buonanno: *A&A*, **372**, 851 (2001)
2. H. Baumgardt, J. Makino: *MNRAS*, **340**, 227 (2003)
3. B. Dauphole, M. Geffert, J. Colin, C. Ducourant, M. Odenkirchen, H. Ticholke: *A&A*, **313**, 119 (1996)
4. G. De Marchi, F. Paresce, L. Pulone: *ApJ*, **530**, 342 (2000)
5. G. De Marchi, L. Pulone, F. Paresce: *A&A*, **449**, 161 (2006)
6. D. Dinescu, T. Girard, W. van Altena: *AJ*, **117**, 1792 (1999)
7. O. Gnedin, J. Ostriker: *ApJ*, **486**, 581 (1997)
8. M. Odenkirchen, P. Brosche, M. Geffert, H. Ticholke: *NewA*, **2**, 477 (1997)
9. E. Vesperini, D. Heggie: *MNRAS*, 289, **898** (1997)